



# Plagiarism Checker X Originality Report

**Similarity Found: 31%**

Date: Thursday, January 23, 2020

Statistics: 806 words Plagiarized / 2576 Total words

Remarks: Medium Plagiarism Detected - Your Document needs Selective Improvement.

---

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/330828527> **New Approach for Referral Preparation to the Health Care Center in Denpasar City and Bogor Regency Article** in **Indian Journal of Public Health Research and Development** · December 2018 DOI: 10.5958/0976-5506.2018.02049.1

CITATIONS 0 READS 393 authors, including: Some of the authors of this publication are also working on these related projects: Big data analysis View project Yuni Rahyani Ministry of Health, Indonesia 9 PUBLICATIONS 1 CITATION SEE PROFILE All content following this page was uploaded by Yuni Rahyani on 12 March 2019. The user has requested enhancement of the downloaded file.

Volume 9 Number 12 December 2018 **New Approach for Referral Preparation to the Health Care Center in Denpasar City and Bogor Regency** Ni Komang Yuni Rahyani 1, Gusti Agung Ayu Budi 2, Wayan Dian Ekayanti 2 1Midwifery Department Polytechnique of Health, Ministry of Health, Diploma 4 Study Program, Jalan Raya Puputan Nomor 11 A Renon Denpasar, Bali, Indonesia, 2Midwifery Department Polytechnique of Health, Ministry of Health, Diploma 3 Study Program, Bogor City, West of Java, Indonesia  
ABSTRACT Knowledge, attitudes and skills of pregnant women to recognize danger signs during pregnancy to postpartum period and newborn baby is a strategic effort to prevent maternal and infant mortality. The aim of the study after guidance at three Health Centers in Denpasar City and Bogor Regency.

The study design was quasi-experimental with nonrandomized pretest-posttest with control group design. The research subjects were maternal and child health books) and control groups (maternal and child health books). The instrument used was a

questionnaire containing the socio demographic characteristics of the subject, knowledge, attitude and observation of skills.

The results showed that there were differences in the scores of knowledge, attitudes and skills in the intervention among groups 1 and 2 ( $p < 0.05$ ). The multivariate test results showed that there guidance intervention, maternal occupation, income and residence ( $p < 0.005$ ). It is necessary to consider the awareness of midwives at the health center to innovate in order to increase alertness and early detection through family-based empowerment.

danger signs guidance, pregnancy, health center, empowerment INTRODUCTION  
According to estimates from world health organizations or the World Health Organization (WHO), as many as 358,000 maternal deaths occur every year (WHO,1 and as many as 88-89% of these deaths can be avoided. The attitudes of health workers (especially midwives and doctors), besides motivation, knowledge of the need for services that forming an important basis for improving maternal and infant health 2,3 .

The maternal mortality rate (MMR) in Indonesia according to data from the Indonesian Demographic and Health Survey (IDHS) in 2012 was 359 per 100,000 live births. The MMR is much higher compared to the 2007 IDHS results of 228 per 100,000 live births. Maternal and newborn health conditions still need serious attention from various parties, not only in Java, but also in Bali.

Thus, basic efforts are carried out by midwives in order to prevent death and illness in mothers and children through efforts to increase the knowledge and awareness of pregnant women and families regarding danger signs to mothers and infants. This effort has a large impact to prevent late referrals, especially the delay in recognizing problems that generally cause delays in making decisions and referring to health facilities 4. to 16.1 per 100,000 live birth but increased back in 2015 and 2016.

Maternal mortality rates in the Denpasar City in 2016 (54 per 100,000 live birth) still lower than the 1400 Indian Journal of Public Health Research & Development, December 2018, Vol. 9, No. 12 in 2016 (100 per 100,000 KH). During 2016 in Denpasar City there were 7 maternal deaths consisting of 5 deaths of pregnant women and 2 during delivery. All mothers die in health facilities (hospitals).

Four maternal deaths in Denpasar City are caused by Non Obstetric Disease, namely 2 people due to heart disorders, 2 people with dengue syock syndrome. The main objective of this research is to compare the skills of pregnant women and their husbands / families regarding danger signs to pregnant women and infants to prevent

delays in obtaining appropriate care between before and after guidance.

In this study, personal guidance methods were used in the two intervention groups and one control group. Guidance media is hand regarding danger signs picture, pocket books and maternal and child health books. We need to know the most appropriate approach to increase knowledge, attitudes and skills of pregnant women and families to prevent delays in recognizing problems and seeking help. MATERIALS AND METHOD The study design was quasi-experimental, pretest- posttest control group design.

Respondents were divided into two groups, namely the control group and the treatment group (1 and 2). The control group (pregnant women in South Denpasar II and Ciomas) were given guidance according to the MCH books at the health center or Puskesmas. Intervention group 1 (Puskesmas I Denpasar Timur and Ciri Mekar) were given guidance and right hand regarding danger signs picture and maternal and child health/MCH books. Intervention group 2 (Puskesmas II Denpasar Barat and Sukaraja). given guidance using pocket books and fans and MCH books.

The sample size in the control and treatment groups will be calculated according to the population of second trimester to third trimester of pregnant women in selected health centers in Denpasar City and Bogor Regency, which amounted to 34 pregnant women in each selected health center or 204 pregnant women in health center. Conducted pretest and posttest for pregnant women regarding danger signs for pregnant women and newborns.

Data collection is carried out for three months (July to September 2017). Pregnant women and husbands who were involved in intervention groups 1 and 2 were given a pretest before guidance. In the control group, pregnant women were given a pretest and were then asked to read maternal and child health books regarding danger signs and preparation for referrals. The time taken from the pretest to posttest is one month.

Guidance in the intervention group and the control group was carried out after the pretest 3 meetings for one hour at each meeting. Guidance is carried out by the MCH management midwives who have been trained in the selected health care. Data analysis was univariate, bivariate and multivariate analysis. Univariate analysis by calculating the mean.

Bivariate analysis to analyze the relationship between independent variables and dependent variables, the intermediate variable with the dependent variable using paired t test and chi square analysis. Multivariate analysis to analyze the relationship between all research variables using multiple regression tests and modeling. RESULTS The

homogeneity test shows if the respondent characteristics are generally homogeneous ( $p > 0.05$ ).

There were differences in the pretest and posttest scores in the intervention group on the score of knowledge, attitudes and skills ( $p < 0.05$ ) except in the control group did not show differences in the skill score ( $p > 0.05$ ). The difference in knowledge, attitudes, and skills scores when compared with intervention group 1 and control group.

In the table 1 below, the results of differences in pretest and posttest scores were displayed on the knowledge, attitudes and skills of pregnant women. Indian Journal of Public Health Research & Development, December 2018, Vol. 9, No. 12 1401 Table 1. Differences in Pretest and Posttest Scores between Intervention and Control Groups (n=204 pregnant women) adjusted variance Knowledge Intervention group 1 24650.75 2.357 0.18 2 24528.00 4.431 0.000 Control 25068.00 2.239 0.025 Attitude Intervention group 1 26258.38 1.858 0.063 2 26451.13 3.034 0.002 Control 25909.38 1.948 0.051 Skill Intervention group 1 26757.13 7.070 0.000 2 26746.63 7.145 0.000 Control 26755.13 1.284 0.199 Source : Primary data analysis (2017) The multivariate test results using general linear model (GLM) analysis showed that there was no maternal age with a knowledge score ( $p > 0.05$ ). There maternal occupational factors with attitude scores ( $p > 0.05$ ).

Consistently intervention, maternal occupation, associated with maternal skills scores regarding danger signs for mothers and newborns baby and neonates ( $p < 0.05$ ). The table 2 below shows the results of the analysis between the intervention, the work of the mother, income and place of residence with the skill score. 1402 Indian Journal of Public Health Research & Development, December 2018, Vol. 9, No. 12 Table 2.

Results of General Linear Model Analysis between Independent Variable, Intermediate and Skill (n=204 pregnant women) in 2017 Skills z p 95% CI AIC Intervention group 7.198 14.083 3.52 8.61 0.000 0.000 10.875 -17.291 7.275 Mother occupation 4.356 2.98 0.003 1.494 -7.219 Family income -7.746 -4.20 0.000 -11.365 -4.127 Residence -8.217 -3.19 0.001 -13.263 -3.171 Husband education: Middle High 1.783 1.719 1.27 0.76 0.204 0.446 -0.967 - 4.532 -2.704 - 6.142 Health assurance -1.166 -0.88 0.380 -3.769 - 1.436 Age 1.200 0.79 0.430 -1.778 - 4.179 Husband occupation -1.376 -0.36 0.721 -8.919 - 6.166 Constanta 13.773 3.65 0.000 6.371 -21.175 Source : Primary data analysis (2017) Discussions In general the characteristics between the intervention groups were not different ( $p > 0.05$ ).

Differences were found from the characteristics of income or socio-economic level, residence, and means of transportation used by subjects to health facilities ( $p < 0.05$ ).

The sociodemographic characteristics of the subject can affect service quality as measured by **knowledge, attitudes and skills** the subject knows and manages the problems faced.

According to the World Health Organization 1, there health facilities, especially to obtain reproductive health distance factors to health facilities, client knowledge, education level and social status economy. Agreeing with the results of a study in Ethiopia by Tesfahun, 5, that maternal health care **cultural beliefs, individual attitudes and** health seeking behavior.

There were differences in the pretest scores of knowledge, attitudes and skills between the intervention ( $p < 0.05$ ), although the results showed that the most The skill score showed that only the control group who compared with intervention group 1 and 2. This was caused by factors of resistance or design invalidity, including: the interaction of the given test, the interaction of the results of the selection with the results and the existence selection with the results and the existence selection bias 6.

The selection bias in terms of the invalidity of interaction factors from the intended test is that there is a weakness of the video media in the intervention group 1 due to the availability of a video watching device at home that is not shared by all subjects in the group, in addition to the obstacles that explain content or content of the video.

Another problem with invalidity is that the characteristics of the subject from the beginning have been different, especially in the intervention group two which shows if the initial results of the pretest are much higher than the other groups 6. pretest and posttest scores according to the characteristics of the transportation facilities and the distance to the health facilities with scores of knowledge, attitudes and skills ( $p > 0.05$ ). Knowledge, attitudes and skills economic factors, values that are believed, and intention to behave.

Different results were obtained by Tesfahun, Indian Journal of Public Health Research & Development, December 2018, Vol. 9, No. 12 1403 5 the utilization of postpartum care are: time constraints, transportation costs, shelter, distance from home to health facilities, previous ANC history and ability health workers provide solutions to problems faced.

The results of multivariate tests using GLM were between intervention, maternal work, family income, and place of residence with a skill score ( $p < 0.05$ ). The results of the study by Amenu, Mulaw, Seyoum & Bayu 7 and Salam, Lassi, Das and Bhutta 8 that positive health complex phenomena related to availability, quality, characteristics of

clients or service users 9 .The success of communication between midwives and clients or satisfaction, increases understanding, increases mutual trust and increases the ability of pregnant women and families in decision making 10 .

Mother's knowledge is not directly affected by the number of ANC visits due / midwives and the limited information provided by midwives regarding danger signs to mothers during pregnancy to postpartum, newborn baby and neonates 11-15 . The husband's education contributes to the participation of husbands in delivering mothers to health facilities and helping to make decisions more precisely and quickly 16 . CONCLUSION The use of media and interesting information delivery techniques are determinants of the success of training.

It is necessary to consider the characteristics of the research subjects, such as the availability of VCD devices, other communication media in the form of mobile phones for intervention groups that receive training using video media. Research Funding : Obtained from Health Polytechnique, Den Pasar Bali. Ethical Clearance: From Faculty of Medicine Udayana University and Sanglah Hospital, no. 1556/ UN.14.2/KEP/2017. in this research and publication. REFERENCES 1.

World Health Organization (WHO),. Social Determinants of sexual and reproductive health. Informing future research and programme implementation. WHO, Geneva: Switzerland. 2010. 2. Hussein, J., Mavalankar, D.V., Sharma, S & D'Ambruoso, L.. A review of health system infection control measures in developing countries: what can be learned to reduce maternal mortality.

Globalization and Health 2011; 7, 14. 3. Hussein,J., Binns, A.McCaw., & Webber, R.. Maternal and perinatal health in Developing Countries.FSC, 2012.; CABI. 4. Mumtaz, Z. & Salway, S..Understanding gendered Pakistan: Moving beyond the autonomy paradigm. Social science and Medicine. 2009; 68, 1349-1356. 5. M. Knowledge, perception and utilization of postnatal care of mothers in Gondar Zuria District, Ethiopia: A Cross-sectional study.

Matern Child Health Journal 2014; (18); 2341 – 2351. 6. Campell,D.T., and Stanley, J.C. Experimental and quasi-experimental designs for research. Rand McNally College Company, Chicago, USA. 1966. 7. Amenu, G., Mulaw, Z., Seyoum, T., and Bayu,H.. Knowledge about danger signs of obstetric complications and associated factors among postnatal mothers of Mechekel District Health centers, east Gojjam Zone, Northwest Ethiopia, 1-7 8. Salam, R.A., Lassi, Z.S., Das, J.K., and Bhutta, Z.A..

Evidence from district evel inputs to improve quality of care for maternal and newborn

health: 2014; 11 (suppl 2): S3 9. Zeleke, D., Nega, A., and Gudha, E. Maternal health care use among married women in Hossaina, Ethiopia. BMC Health Serv Res. 2015; 15: 36 10. Glanz, K., Rimer, B.K., and Viswanath, K.. Health behavior and health education. Theory, Research, and Practice. 4 th ed. Jossey and Bass; United States of America. 2008; pp 45-65. 11. Kabakyenga, J.K.,

Ostergren, P.O., Turyakira, E, and 1404 Indian Journal of Public Health Research & Development, December 2018, Vol. 9, No. 12 Reprod Health. 12. Birawida, A.B., Selomo, M., Mallongi, A. Potential hazards from hygiene, sanitation and bacterium of (and food safety perspective), IOP Conference Series: Earth and Environmental Science. 2018. 13.

Sriasih, N.G.K., Ariyani, N.W., Mauliku, J., Budiani, N.N., Mallongi, A. Frangipani aromatherapy oil perineum ruptur spontan at the labo. Indian Journal of Public Health Research & Development. 2018. 14. Merta, I.W., Sudarmanto, I.G., Sri Dhyana Putri, I.G.A., Arjani, I.A.M.S., Mallongi, A., Working position improvement by adding supporting tool reduced subjective complains and increase productivity of weavers in tenganan village karangasem regency.

Indian Journal of Public Health Research & Development. 2018. Indian Journal of Public Health Research and Development 16. Hoque, M., and Hoque, M.E.. Knowledge of Danger Signs for Major Obstetric Complications Among Pregnant KwaZulu-Natal Women. Implications for Health. 2011; Desember. 4. View publication stats View publication stats

#### INTERNET SOURCES:

-----  
14% -

[https://www.researchgate.net/publication/330828527\\_New\\_Approach\\_for\\_Referral\\_Preparation\\_to\\_the\\_Health\\_Care\\_Center\\_in\\_Denpasar\\_City\\_and\\_Bogor\\_Regency](https://www.researchgate.net/publication/330828527_New_Approach_for_Referral_Preparation_to_the_Health_Care_Center_in_Denpasar_City_and_Bogor_Regency)

<1% -

[https://www.researchgate.net/publication/7471566\\_Implementation\\_of\\_a\\_SNAP\\_intervention\\_in\\_two\\_divisions\\_of\\_general\\_practice\\_A\\_feasibility\\_study](https://www.researchgate.net/publication/7471566_Implementation_of_a_SNAP_intervention_in_two_divisions_of_general_practice_A_feasibility_study)

1% - [https://www.researchgate.net/profile/Yuni\\_Rahyani](https://www.researchgate.net/profile/Yuni_Rahyani)

<1% - <http://www.ijikm.org/Volume6/IJIKMv6p151-161Levy553.pdf>

<1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4390384/>

<1% - <https://www.sciencedirect.com/science/article/pii/S026069171100308X>

<1% - <https://www.who.int/hiv/en/>

1% -

<http://news.unair.ac.id/en/2018/08/07/siaga-plus-application-supports-pregnant-mother-health-indonesia/>

<1% -

<https://health-policy-systems.biomedcentral.com/articles/10.1186/s12961-017-0213-9>

<1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5001799/>

<1% -

[https://www.researchgate.net/publication/323416883\\_Out-of-pocket\\_expenditure\\_on\\_maternity\\_care\\_for\\_hospital\\_births\\_in\\_Uttar\\_Pradesh\\_India](https://www.researchgate.net/publication/323416883_Out-of-pocket_expenditure_on_maternity_care_for_hospital_births_in_Uttar_Pradesh_India)

<1% - <https://www.liebertpub.com/doi/abs/10.1089/jwh.2004.13.713>

<1% -

[https://www.popline.org/apachesolr-angularjs-search/im\\_field\\_document\\_major\\_keywords%3A55318%20OR%20im\\_field\\_document\\_minor\\_keywords%3A55318](https://www.popline.org/apachesolr-angularjs-search/im_field_document_major_keywords%3A55318%20OR%20im_field_document_minor_keywords%3A55318)

<1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3170621/>

<1% - <https://www.arcjournals.org/pdfs/ajphcm/v2-i3/1.pdf>

<1% - <https://dergipark.org.tr/en/download/article-file/806188>

<1% -

<https://www.engineeringintro.com/statistics/introduction-statistics/univariate-bivariate-and-multivariate-data/>

<1% - <https://www.petropedia.com/definition/439/bivariate-analysis>

<1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3050606/>

<1% - <https://www.sciencedirect.com/science/article/pii/S0168010216301845>

<1% -

<https://www.slideshare.net/AWEYMULYANA1/effects-of-multimedia-instructional-material-on-students-learning>

1% -

<https://www.openaccessjournal.com/journal/81/Indian-journal-of-public-health-research-and-development>

<1% -

<https://medworm.com/rss/medicalfeeds/source/Western+Journal+of+Nursing+Research.xml>

<1% - <https://www.frontiersin.org/articles/10.3389/fpsyg.2017.01179/full>

<1% - <https://bmcpublichealth.biomedcentral.com/articles/10.1186/1471-2458-11-403>

<1% - [https://link.springer.com/chapter/10.1007/978-3-319-73820-8\\_13](https://link.springer.com/chapter/10.1007/978-3-319-73820-8_13)

<1% - <https://www.sciencedirect.com/science/article/pii/S0167945715300701>

<1% -

<https://wattsupwiththat.com/2017/07/23/tesla-battery-subsidy-and-sustainability-fantasies/>

<1% - <https://cis.org/Report/Welfare-Use-Immigrant-Households-Children>

<1% -

<http://repository.unair.ac.id/88761/1/6%20Concentration%20of%20Benzene%20Exposure%20in%20a%20Laboratory%20of%20Oil%20Processing%20Industry.pdf>

<1% - <https://www.sciencedirect.com/science/article/pii/S0266613813000119>



<1% -

<http://www.e-urol-sci.com/article.asp?issn=1879-5226;year=2019;volume=30;issue=5;spage=211;epage=215;aulast=Gede;type=3>

1% - <https://www.popline.org/node/210525>

1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3907541/>

1% - <https://staffprofiles.bournemouth.ac.uk/display/journal-article/34503>

<1% - <https://www.sciencedirect.com/science/article/pii/S0010782414003266>

1% -

[https://www.academia.edu/13700372/Knowledge\\_Perception\\_and\\_Utilization\\_of\\_Postnatal\\_Care\\_of\\_Mothers\\_in\\_Gondar\\_Zuria\\_District\\_Ethiopia\\_A\\_Cross-Sectional\\_Study](https://www.academia.edu/13700372/Knowledge_Perception_and_Utilization_of_Postnatal_Care_of_Mothers_in_Gondar_Zuria_District_Ethiopia_A_Cross-Sectional_Study)

1% -

<https://paperity.org/p/76387192/knowledge-of-obstetric-danger-signs-and-associated-factors-among-pregnant-women-in-erer>

1% - <http://scholar.google.co.id/citations?user=dr2YTzzNtxwC&hl=id>

1% - <http://www.poltekkes-denpasar.ac.id/produk/>

<1% -

[https://www.researchgate.net/journal/0976-0245\\_Indian\\_Journal\\_of\\_Public\\_Health\\_Research\\_and\\_Development](https://www.researchgate.net/journal/0976-0245_Indian_Journal_of_Public_Health_Research_and_Development)

1% - <https://academic.oup.com/her/article/29/6/1041/2804357>

1% -

[http://s3-us-east-2.amazonaws.com/spectralevolution/assets/20171103185821/Siltstone\\_s\\_Shale.pdf](http://s3-us-east-2.amazonaws.com/spectralevolution/assets/20171103185821/Siltstone_s_Shale.pdf)