

Plagiarism Checker X Originality Report

Similarity Found: 3%

Date: Kamis, Maret 19, 2020 Statistics: 91 words Plagiarized / 3441 Total words Remarks: Low Plagiarism Detected - Your Document needs Optional Improvement.

Factors Related to Dental Caries on Elementary School Students in Badung Regency – Indonesia Authors Asep Arifin Senjaya1*, Ni Made Sirat2, I Nyoman Wirata3 1,2,3Department of Dental Health, Health Polytechnic of Health Ministry Denpasar, Indonesia *Corresponding Author Asep Arifin Senjaya Abstract Caries, dental and oral disease affects 90% of school-age children. Badung regency occupied fourth position of the populations with dental problem in past 12 months. The data shows only 3.6% of Badung population who behaved properly in brushing. Badung regency economically has the highest local revenues in Bali.

The objective of this research is to understand the factors associated with dental caries in primary school students in Badung. This research type is quantitative- qualitative, conducted from July to September 2016. The sample was 252 students, selected by multistage random sampling. The result showed 71% of students belong to very high, high, and moderate caries; 64.8% of students belongs to low sosio-economic category and 89.7% of students need guidance in brushing.

The results show there is no relationship between the variables of socio-economic category with a classification of dental caries proven by sig value results in 0.552. There is a relationship between variables of behavior criteria in brushing with dental caries classification with sig. value of 0.000. There is a relationship between school dental health (UKGS) with the classification of dental caries with sig. value of 0.001.

Multivariate analysis shows sub variable UKGS category 1 and sub variable category 2 are associated with dental caries. Keywords: Behaviour, children, dental caries, factors; related. care and parental knowledge4. Parents need to Introduction _observe and regularly record changes occurring at _ _The degree of health of a person or community

is _the child's oral cavity in anticipation of early dental _ _influenced by four factors: environment, attitudes, _disorders and abnormalities, such as dental caries.5 _ _hereditary, and health care1.

There are three factors _The average caries experience (DMF-T) of 12 years _ _affecting one's behaviour, predisposition factors, _old children ranged from 2.4. The DMF-T index _ _supporting factors, and reinforcing factors2. Oral _according to WHO for school-age children up to 12 _ _health is a fundamental part of general health and _years is = 36.

School-age children are one of the __welfare of life. Oral health disorder will have an _groups that are vulnerable to caries, as they __impact on one's performance.3 _generally still have a lack of knowledge and __Dental caries is a dental and oral disease that is _behavior that is lacking in dental caries. ___widely suffered by 90% of school-age children.

_There has been an increase in the prevalence of _ _Caries is also the cause of primary pathology of _active dental caries in Indonesian population _ _decayed tooth that occur due to the lack of child _compared to 2007 years ago, namely from 43.4 in _ _ _ _ _ _

2007 to 53.2% in 2013. More than 93 million Indonesian people suffer from active caries.

People with dental and oral problems in past 12 months in Badung Regency is at the top four in the province of Bali, which is 31.6%7. Plaque that is firmly attached to the surface of the tooth and gums may cause disease in the hard tooth tissue and its supporting tissues8. Riskesdas Data 2013 shows the percentage of the population aged 10 and above who brush their teeth every day in Badung Regency was 94.2% and only 3.6% that behaved properly in brushing 7.Parents ' participation is indispensable in guiding, caring, understanding, reminding, and providing facilities for children to keep their teeth clean 9 Parents have an enormous influence on child behavior, as parents are the first figure to their children. Parents are obligated to give and teach positive things as well as compassion.

9 Dental health services in Health Centre (Puskesmas) can be implemented in and outside the building such as School Dental Health (UKGS) and in Posyandu (Integrated Health Post)10. School Dental Health consists of three stages, namely UKGS phase I, UKGS phase II, and UKGS phase III11. Badung Regency has the largest economic scale in Bali, in terms of regional revenues and expenditure (APBD).

This can be understood because the Badung Regency has the famous tourist destinations 12. The revenue distribution of Bali province in 2013 shows 40% middle-ranking society covers 15.12% of Badung Regency. Badung Regency is second population with low income after the city of Denpasar 13.

Furthermore, the problem can be formulated with "What is the relation of factors with dental caries in elementary school students in Badung Regency in 2016?" The purpose is to understand the factors related to dental caries of elementary school students in Badung Regency in 2016. Method This research uses quantitative and qualitative method.

The research took place in elementary _ schools (SD) class six, in six sub-districts in Badung Regency, namely: SDN 3 Kuta; SDN 11 Jimbaran; SDN 1 Kerobokan; SDN 1 Pangsan; SDN 2 medium; SDN 3 Sempidi. Data collection and information was done from July to September 2016. The research population is all elementary school students and their parents in Badung Regency.

Large sample research is taken with a large formula of cross sectional study14. In this study the number of samples was reproduced, taken 42 students in each SD, so the total number of 252 students and 252 parents, taken with multistage random sampling. The

FGD informant was from the representatives of the parents, each of the elementary school parents (mothers) of students are ten in number.

The in-depth interview informant was a teacher in each of the elected primary schools. Dental caries data is measured by direct examination, tooth brushing with interview and observation, socio-economics using questionnaire, UKGS with interview. While the perception of parents and teachers about dental caries was acquired through FGD and in-depth interview.

Quantitative data analysis was done using univariate, frequency and percentage. Bivariate analysis with cross-tabulation and correlation test Spearmen in multivariate analyses used multi-regression logistics. Qualitative information obtained was used to complement quantitative discussions.

Result and Discussion School-age children are one of the groups vulnerable to caries, as they still have less knowledge and behaviour against dental caries 15. Table 1 shows the frequency distribution of students based on caries they suffer from. Table 1 Student frequency distribution based on dental caries classification Classification of _f (%) _ _ Dental Caries _ _ _ Very high _74 (29,4) _ _high _53 (21,0) _ _moderate _52 (20,6) _ _low _20 (7,9) _ _ Very low _53 (21,0) _ _Total _252(100) _ _

Table 2 shows 164 students (64.8%) included in the low socioeconomic category (average income under Rp. 1,500,000.- / month) and moderate (average income from Rp Rp 1,500,000. - to Rp. 2,500,000.-

/ month), the remaining 34.9% is categorized as high and very high socioeconomic. Based on this data it can be said that most students are poor families. The most influential factor on health according to 1 is the environment. Within these environmental factors, the economy is a part of it. The environment, including socioeconomic, has the greatest role in health.

Many studies show that the prevalence of caries is higher in children who come from low socioeconomic status. _ Table 2 Frequency distribution by socio-economic category Socio-economic category _f (%) _ _Low _51 (20,0) _ _Moderate _113 (44,8) _ _High _56 (22,2) _ _Very high _32 (12,7) _ _Total _252(100) _ _ Table 3 shows 226 students (89.7%) need guidance in brushing their teeth.

Only 2 students had excellent tooth brushing behaviour and 3 students had good tooth brushing behavior. This data is in accordance with the results of Basic Health Research in 20137, where in the population in Badung Regency 94.2% brushed teeth every day, but only 3.6% behaved correctly in brushing.

Table 3 Frequency distribution by socio-economic category No.

_Age _Intervention Group _Sample Control Group _Total _ _ _ _ Group _f _% _f _% _f _ _% _ _1. _8 _64 _51.6 _60 _48,4 _124 _ _50 _ _2. _8,5 _1 _8 _- _- _1 _ _0,4 _ _3. _9 _53 _42.7 _63 _50,8 _116 _ _46.8 _ _4. _10 _6 _4,9 _1 _0.8 _7 _ _2.8 _ _ _Total _124 _100 _124 _100 _248 _ _100 _ _

All informants said that if a tooth has a cavity, it must be treated immediately, that is, it is patched to the dentist, not to cause pain / swelling. But if it cannot be treated it should be removed and made denture.

The role of parents, especially a mother in maintaining dental health is very important in underlying the formation of behaviors that support children's dental hygiene so that children's dental health can be maintained properly 16. Most of the informants said that they should remind their children to brush their teeth. A small number of informants said not always, some informants even said that it was rare and some said that they did not remind again with the reason that they were bored, because as if the words of their parents were ignored, the children preferred to play mobile phones / games rather than being told to brush their teeth.

But actually, all the informants agreed that parents should not be bored and must continue to remind their children to brush their teeth until the child has a habit of being independent in brushing teeth. _ The study found inactive UKGS in SDN 1 Pangsan, SDN 3 Sempidi in UKGS Phase I, and four other SDNs namely: SDN 11 Jimbaran, SDN 3 Kuta, SDN 1 Kerobokan, and SDN 2 Sedang, the UKGS was less active.

The results of this study are similar to the conclusions of a study conducted by 17 which stated that the UKGS program in the Polonia Puskesmas's work area was not maximal. Dental health services can be carried out in or outside Puskesmas such as the School Dental Health Efforts (UKGS) at schools and at Posyandu. Puskesmas have an important role in the implementation of the UKGS program11.

The implementation of the UKGS involved a number of staff, from Puskesmas namely dentists and dental nurses, while the implementers from the schools were health sports teachers. The attitude of the health sports teacher towards the UKGS program is very influential on the changes in students' attitudes to oral health and the success of this program10.

The cross table between the socioeconomic categories and the classification of dental caries

relationship between the two variables with the strength of the weak relationship in the opposite direction. The results of this study are different from the results of research conducted previously where students in primary schools who did not carry out UKGS program have a higher risk of caries exposure compared to students in primary schools who receive the UKGS program20.

Table 5 Cross tabulation criteria with dental caries classification Classification _of _ _ Socio-economic category _ Total (%) _ _dental caries _ _Need guidance _Moderately _ Good n(%) _ Very good _ _ _ _ _ n(%) _ _ n(%) _ _ _ Very high _ _ 73 (98,6) _ 1 _ (1,4) _ 0 (0) _ 0 (0) _ 74 (100) _ _high _ _ 49 (92,5) _ 4 _ (7,5) _ 0 (0) _ 0 (0) _ 53 (100) _ _ moderate _ _ 43 (82,7) _ 6 (11,5) _ 2 (3,8) _ 1 (1,9) _ 52 (100) _ _low _ _ 16 (80) _ 3 _ (15) _ 1 (5) _ 0 (0) _ 20 (100) _ _ Very low _ _ 45 (89,9) _ 7 (13,2) _ 0 (0) _ 1 (1,9) _ 53 (100) _ _ Total _ _ 226 (89,7) _ 21 (8,3) _ 3 (1,2) _ 2 (0,8) _ 252 (100) _ _ The conclusions of (Anitasari and Rahayu)18 states that there is a correlation between the frequency of tooth brushing and the level of dental and oral hygiene. Table 6 shows the better implementation of the UKGS, the higher the caries rate.

Spearmen correlation test between UKGS category variables and dental caries classification obtained sig. 0.001 and the correlation coefficient (r) -0.252. This means that there is a Table 6 Cross tabulation category UKGS with dental caries classification _Classification of _ _ UKGS category _ _ Total (%) _ _ _ dental caries _No UKGS n(%) _ UKGS Inactive n(%) _UKGS Phase I n(%) _ _ _ Very high _7 _(9,5) _47 (63,5) _20 _(27,0) _74 (100) _ _ _ High _10 _(18,8) _31 (58,5) _12 _(22,7) _53 (100) _ _ _ Moderate _9 (17,3) _34 (65,4) _9 (17,3) _52 (100) _ _ _ Low _3 (15) _17 (85) _0 (0) _20 (100) _ _ _ Very low _13 _(24,5) _39 (73,6) _1 _(1,9) _53 (100) _ _ _ Total _42 _(16,6) _168 (66,8) _42(16,6) _252 (100) _ _ _ _

Table 7 shows SDN 3 Sempidi (UKGS Phase I category) having the most students who need guidance on brushing and none of their students behave well and very well.

This factor may have contributed to the high caries in SDN 3 Sempidi. Though they are in the UKGS Phase I category, their teeth brushing behaviour is bad. Unlike the case with SDN 1 Pangsan, although in SDN 1 Pangsan there was no UKGS, but his teeth-brushing behaviour was generally the best, there were found two students with excellent tooth-brushing behaviour. In (Suryani)20 shows that the scope of UKGS program had not met the target.

The supervision and control of UKGS program does not yet have clear indicators. Supervision is only carried Table 7 Elementary tabulation with dental behaviour _ out by a small proportion of Puskesmas. Lack of guidance and supervision at the level of district and unclear UKGS structure at the provincial level.

This is also possible in Badung District, which has an impact on outputs that are not as expected. In addition, caries etiology is very much, not only from UKGS program. The internal factors are directly related, consisting of four factors, namely the host (teeth and saliva), the causative agent of the disease (microorganisms in plaque), the substrate (carbohydrate), and the time.

External factors include factors such as age, ethnicity, culture, social population, awareness, attitudes, and behaviour of individuals towards dental health.

Elementary School __Dental Brushing Behavior __Total ___Need __Moderate _Good __Very good _ _ _ guidance. _ _ _ _ SDN 11 Jinbaran _39 _ 3 _0 _0 _42 _ SDN 3 Kuta _37 _ _5 _0 _0 _42 _ SDN 1 Kerobokan _39 _ 3 _0 _0 _42 _ SDN 3 Sempidi _41 _ _1 _0 _0 _42 _ SDN 2 Sedang _37 _ _4 _1 _0 _42 _ SDN 1 Pangsan _33 _ _5 _2 _2 _42 _ _

Table 8 shows logistic regression multivariate test results of UKGS variable and tooth brushing behavior variable that were simultaneously tested with dental caries variable, showing the results of the UKGS sub variable category 1 (UKGS not active / no UKGS) (sig. 0.001) and the UKGS sub variable category 2 (UKGS less active / there is a UKGS but not Phase I) (sig. 0,000).

The results of this test can be interpreted if the UKGS variable and the variable brushing behaviour are simultaneously tested with dental caries variable, then UKGS _ sub variable category 1 (UKGS is inactive / no UKGS) and the UKGS sub variable category 2 (UKGS is less active / there is UKGS but not Phase associated with dental caries.

This can be understood even though SDN 3 Sempidi is included in the UKGS category 3 (UKGS Phase I), but because almost all respondents way of brushing still need guidance and other factors not examined in this study, which may be a cause of high caries in SDN 3 Sempidi So that in this study UKGS seems not useful in preventing dental caries.

Conclusions Most students of SDN in Badung Regency have dental caries with classification of high and very high caries. They come from moderate and low socio-economics and need guidance of brushing their teeth.

Most of the UKGS at SDN in Badung Regency are inactive categories. Regarding socie-economic and dental caries on students at SDN Kabupaten Badung, there is no relation between those variables but there is a relationship between brushing and dental caries on students at SDN Kabupaten Badung as well as there is a relationship between UKGS and dental caries on students at SDN Kabupaten Badung.

Either inactive or less active UKGS, they have relation to dental caries on students at SDN Kabupaten Badung. Qualitative study concluded that parents of students and sports teachers have a good knowledge and attitude about dental caries.

Acknowledgement Obtained from the Committee of Respondents agreement. Source of Funding: Ministry of Health Republic Indonesia. References Notoatmodjo, S.

Education and health behavior. Jakarta: Rineka Cipta. 2003 Sriyono, NW. Preventing dental and oral diseases in order to improve quality of life, speeches of the inaugural Department of the Faculty of Dentistry at Universitas Gadjah Mada Yogyakarta. 2009. Putri MH, Herijulianti E, Nurjanah N. The science of disease prevention hard Network and dental support network, Jakarta: EGC. 2011.

Itjingningsih W H. Anatomi Gigi. Jakarta: EGC. 1991. World Health Organization (WHO). Oral HealthInformationSystems.2003. Available at: www.who.int/oral health/action/information. Accessed January 19, 2016. _ Ministry of Health (Depkes) RI. How to work with dental and oral health care in Puskesmas. Jakarta: Directorate of Dental Hygiene. 1995. Suciari A, Arief YS, Rachmawati PD. Parents role in guiding teeth brushing with the genesis of preschool children's dental caries. 2015. Available at: journal.unair.ac.id/download-fullpapers-pmnj5ab2f12.

Retrieved: 18 January 2016. Puspitoningsih N, Safitri W, Istiningtyas A. Mother's perception of dental caries in preschool-age children in TK Darma Women's District Kemusu Boyolali. 2015. Available at www.digilib.stikeskusumahusada.ac.id. Retrieved: 18 January 2016. Ministry of Health () RI. Dental and oral helath care guidelines (UKGS). Jakarta: Directorate General of Bina Kesehatan Health. 2012.

Septiasih. Badung has the largest economic scale in Bali. 2015. Available at https://www.selasar.com. Retrieved 20 January 2016. Central statistic Agency (BPS). Bali in numbers 2014. Available at: Bali bps.go.id. Accessed in 18 January 2016. Murti B. Principles and methods of epidemiology research. Yogyakarta: Gadjah Mada University

Press. 1997 World Health Organization (WHO). Oral Health Surveys Basic Methods – 5th edition. 2013. Available at: www.who.int/oralhealth/publication.

Retrieved 20 January 2016. Suwelo I S. Dental caries in children with a variety of etiology factors: study on preschool age children. Jakarta: EGC. 1992. Tunggadewi G. Analysis of the implementation of Dental Health Business Program (UKGS) in the region of Polonia district of Medan Polonia in 2014. Thesis. 2015. Available at:www.digilib.usu.ac.id. Accessed: 18 Jan 2016.

Ngangtung R, Pangemanan HC, Gunawan PN.

Influence of the social eonomic level of parents to children's caries in the TK Hang Tuah Bitung. 2015. Available at: E-Dental Journal (eG) Volume 3, Number 2, July-December 2,015,542. Accessed 18 January 2016. Susi, Bachtiar H, Azmi U. Relationship of the elderly socio-economic status with caries on the eldest teeth of children aged 4 and 5 years. Andalas Medical Magazine No. 1. Vol.36.

January 2012. Available at: jurnalmka,fk,unand.ac.id/index.php/article. Accessed 18 January 2016. Anitasari S, Rahayu N E. Relations frequency brushing with dental hygiene and mouth level students of public elementary school in District Palaran Samarinda Municipality of East Kalimantan Province. Dentistry magazine. (Dental Journal), Vol. 38. 2 April – June 2005:88 – 90. Available at: journal.unair.ac.id. Retrieved 18 October 2016.

Kamayanti I A, Tauchid S N, Harahap N N. Risk difference in the occurrence of new caries in children aged 12 years of elementary school students of UKGS and SD non UKGS in the district of Cilandak South Jakarta in 2011. Research report. Available at: poltekkesjakarta.ac.id/file/dokumen. Retrieved 20 January 2016. Suryani Y.

Analysis of the management of dental health business Program at the Healthcenter of the school of Jakarta Barat year 2002. Thesis. Available at: http://lib.ui.ac.id/bo/uibo. Accessed 19 January 2016.

INTERNET SOURCES:

<1% -

https://www.researchgate.net/publication/233880063_Association_between_Obesity_and _Dental_Caries_in_Chinese_Children

<1% -

https://www.researchgate.net/publication/235796524_Early_Childhood_Caries_and_Body _Mass_Index_in_Young_Children_from_Low_Income_Families

- <1% https://e-journal.unair.ac.id/MKG/article/download/3571/2586
- <1% https://www.journalcra.com/archive/201611
- <1% https://www.freekidstories.org/parents/category/emotional%20intelligence/2
- <1% https://www.unud.ac.id/en/daftar-ta.html
- < 1% https://www.science.gov/topicpages/p/preferred+teaching+techniques.html
- <1% https://id.scribd.com/doc/294910285/Statistics-in-Action
- <1% http://www.fao.org/3/y5825e/y5825e07.htm
- <1% https://bmcoralhealth.biomedcentral.com/articles/10.1186/s12903-016-0318-5
- <1% -

https://www.researchgate.net/publication/321045784_A_Cross-Sectional_Study_of_Nutrient_Intake_and_Health_Status_among_Older_Adults_in_Yogyakarta_Indonesia <1% -

https://www.childrenshealthfund.org/wp-content/uploads/2017/02/HBL-Dental.pdf 1% - https://pubs.opengroup.org/architecture/archimate3-doc/apdxb.html <1% -

https://es.scribd.com/document/393921387/ICGH-2017-Proceeding-Update-ISBN

- <1% https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3255964/
- <1% https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2566974/
- <1% https://zombiedoc.com/proceeding-isbn-978-602-73585-0-8.html
- <1% -

https://www.researchgate.net/publication/282604453_Assessing_Learning_Barriers_amo ng_Dental_and_Nursing_Undergraduates_A_Qualitative_Study_Students'_Perspective 1% - https://db0nus869y26v.cloudfront.net/en/Assistant_Bishop_of_Newcastle